



JAEDYN HINDS

ACL Readiness Screen · SDSC

Date: 2026-06-11


Age: 15 yrs

Platform: VALD ForceDecks



JUMP BALANCE — LEFT VS RIGHT

A jump has three phases — she **loads & brakes** on the way down, **pushes off**, then **lands**. We measure how evenly her legs share the work in each, plus a single-leg test. Bars show the left-vs-right gap (shorter = more balanced; target under 10%).




What this means. On one leg, her **right leg** produces noticeably less power and control than the other — a meaningful left-vs-right difference. On the pitch that shows up in her first-step acceleration off that leg, her balance landing from headers and 50/50 challenges, and how she holds a cut. A gap like this is one of the most established, **modifiable** risk factors for a non-contact knee injury — far better caught now, on a force plate, than after an injury. The good news: it's very trainable.

This is a movement **screen**, not a medical diagnosis. The next pages explain how an ACL injury happens, what this means for her game, her numbers, and the plan. Captured on VALD ForceDecks (dual force plates). © Catalyst Performance.

JAEDYN — UNDERSTANDING THE RESULT

HOW A NON-CONTACT ACL INJURY HAPPENS



The **ACL** is the main ligament that stabilises the knee — it stops the shin sliding forward and the knee twisting inward. About **70% of ACL tears are non-contact**: they happen in a split second when an athlete **decelerates, lands, or cuts** and **the knee caves inward** (the position on the right).

A leg that's **weaker or less controlled is more likely to collapse into that position** — and less able to absorb the load — so the ligament takes the strain. **That's why a left-vs-right asymmetry matters: it flags the leg least able to handle the exact forces that tear an ACL.** For jaedyn, that's her **right leg** — the side least able to control a one-leg landing or cut, which is when the ACL is most exposed.

WHAT THIS MEANS FOR JAEDYN'S GAME

ON THE PITCH

- **First-step acceleration** — Off her right leg she drives with less power — that costs her a step.
- **Landing from headers & 50/50s** — Less single-leg control means less stability on impact.
- **Holding a cut** — Planting and changing direction on the weaker side is harder to control.

WHY IT MATTERS — THE EVIDENCE

- Female soccer players are **4–6× more likely** to tear an ACL than boys.
- About **70% are non-contact** — on cuts, decelerations and landings, the exact movements measured here.
- A left-vs-right gap over **10–15%** is a recognised, **modifiable** risk factor and a return-to-play standard.
- Soccer-specific neuromuscular training (FIFA 11+) cuts ACL injuries **30–50%** — and works best in youth.

Refs: Arendt & Dick 1995; Grindem 2016; Bishop 2018; Sadoghi 2012 (meta-analysis).

MORE THAN INJURY — PERFORMANCE

Symmetry isn't only protective — it's faster. The same balance that guards the knee also drives her game:

- **Acceleration & cutting** — Even drive off both legs means a quicker first step and sharper change of direction.
- **Power transfer** — Her jump of **23.4 cm** (Average for her age) is her explosive base — balanced legs put it into sprints, jumps and shots.
- **Durability** — When one leg isn't overworking, she fatigues less and stays available all season.

JAEDYN — HER NUMBERS & PLAN

HOW JAEDYN'S NUMBERS COMPARE

MEASURE	JAEDYN	TARGET / REFERENCE	STATUS
Single-leg jump symmetry (L vs R)	16% gap	<10% gap (≥90% symmetry)	Follow-up
Push-off balance (take-off)	2% gap	<10% gap	On target
Load / brake balance (decelerating)	12% gap	<10% gap	Monitor
Landing balance (impact)	19% gap	<10% gap	Monitor
Counter-movement jump height	23.4 cm	Average vs girls age 15	Baseline

Jump Height vs Girls Her Age (15) — jaedyn 23.4cm · Average



Symmetry targets follow return-to-sport standards (≥90% limb symmetry). The jump-height benchmark is an approximate developmental reference for female youth, not a clinical percentile.

WHAT TO WORK ON — JAEDYN'S TAKE-HOME

- 1 Eccentric single-leg strength — 3-second-lowering split squats and single-leg RDLs — both sides, extra reps on the weaker leg. The braking phase is when the knee is under the most load.
- 2 Decelerate-and-stick — Jog, plant on one leg and freeze for 2 seconds — it trains each leg to absorb force evenly.

Coach-led general guidance from today's screen — a starting point, not a prescription. The complimentary follow-up adds the single-leg test and turns this into a personalized, supervised plan with a re-test to show progress.

HOW WE CAN HELP — NEXT STEPS

FREE RESULTS CALL
Complimentary · 15 min

Walk through jaedyn's results with our staff and map out what to do next.

IN-DEPTH EVALUATION
\$169 · 1 hour

A fuller VALD force-plate battery — more movements tested to pinpoint the cause and build her plan.

COMPLETE ASSESSMENT
\$349 · 2 hours

The full VALD evaluation **plus** hands-on flexibility & mobility screening with our ATC on the table.

BOOK JAEDYN'S SESSION

Recommended given today's result — reply to this email, text us, or talk to your coach — we'll get her started.

How this was measured. Captured on **VALD ForceDecks** — dual force plates sampling **1,000x/second**, the same system used by professional and national teams. Every value is the **average of 3 maximal jumps** for reliability, and compared against published youth-athlete reference ranges. Prepared by the Catalyst Performance team.

Important. Wellness **screen**, not a medical diagnosis or injury prediction — asymmetry is one risk indicator, not a guarantee either way. Results vary day to day; any pain or prior injury should be seen by a physician/PT. © Catalyst Performance · catalystperformancesd.com